

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP PROMAX® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup PROMAX Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

DOC ID 521252

ACCEPTED
FOR REGISTRATION

FEB 4, 2010

New York State Department
of Environmental Conservation
Division of Solid & Hazardous Materials
Pesticide Product Registration

EPA Reg. No. 524-579



Roundup PROMAX is a registered trademark
of Monsanto Technology LLC.

For Foliar and Broadcast Treatment of Japanese Knotweed & Oriental Bittersweet

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of the herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING AND APPLICATION INSTRUCTIONS" sections of the label booklet for Roundup PROMAX Herbicide for essential product performance information. See the "WEEDS CONTROLLED" section of the label booklet for Roundup PROMAX Herbicide for optimum stage of treatment of weed for best results.

JAPANESE KNOTWEED

For control of Japanese knotweed (*Polygonum cuspidatum*), this product may be applied as a 1.5% v/v spray-to-wet solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

For broadcast applications, apply 2.75 quarts of this product in 3 to 40 gallons per acre as a broadcast treatment.

Late summer or fall application is ideal, while leaves are still green and after fruit formation.

ORIENTAL BITTERSWEET

For control of Oriental bittersweet (*Celastrus orbiculatus*), this product may be applied as a 1.5% v/v spray-to-wet solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

For broadcast applications, apply 2 quarts of this product in 3 to 40 gallons per acre as a broadcast treatment.

Late summer or fall application is ideal, while leaves are still green and after fruit formation.

WARRANTY AND LIABILITY

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup PROMAX herbicide before buying or using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

MONSANTO COMPANY
800 N. Lindbergh Blvd.
St. Louis, Missouri, 63167 U.S.A.
©2009

7/1/2009

63028H1-25