Water Soluble Granule

Roundup QuikPRO™ Herbicide

Roundup QuikPRO herbicide is a fast-acting, non-selective professional herbicide for use in non-crop areas and industrial sites.

Roundup QuikPRO herbicide
Dead Weeds — Quick



ACCEPTED
FOR REGISTRATION

10/20/11

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc. ID: 528652

Keep Out of Reach of Children.

CAUTION! See back panel booklet for additional precautionary statements.

Manténgase fuera del alcance de los niños.

¡PRECAUCION! Vea trasero el librete del panel para las declaraciones preventivas adicionales.

ACTIVE INGREDIENTS:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of its ammonium salt	73.3%
Diquat dibromide [6,7-dihydrodipyrido (1,2-a;2',1'-c) pyrazinedilium dibromide]	2.9%
OTHER INGREDIENTS:	23.8%
	100.0%

*Equivalent to 66.6% of the acid, glyphosate, 1.0 pound of Roundup QuikPRO herbicide contains 0.73 pound of the ammonium salt of glyphosate and 0.03 pound of the dibromide salt of diquat.

See label booklet for precautions and instruction in English and Spanish. Vea el librete de la etiqueta para las precauciones y la instrucción en inglés y español.

NET WT 6.8 LBS

Container filled by weight, not volume. 98005J7-9/53

FRONT COVER
(PAGE 1)



FPO
UPC: No barcode in IRD

BACK COVER
(LAST PAGE)

Roundup QuikPRO™ Herbicide

Roundup QuikPRO herbicide is a fast-acting, non-selective professional herbicide for use in non-crop areas and industrial sites.

Complete Directions for Use

AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE, STEMS, EXPOSED NON-WOODY ROOTS, OR DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

EPA Reg. No. 524-535 2811-1

Read the entire label before using this product.

Use only according to label instructions.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

1.0 INGREDIENTS

ACTIVE INGREDIENTS:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its ammonium salt	73.3%
Diquat dibromide [6,7-dihydrodipyrido (1,2-a:2',1'-c) pyrazinediium dibromide]	2.9%
OTHER INGREDIENTS	23.8%
	100.0%

*Equivalent to 66.6% of the acid, glyphosate.

1.0 pound contains 0.73 pound of the ammonium salt of glyphosate and 0.03 pound of the dibromide salt of diquat.

This product is protected by U.S. Patent No. 7,008,904. Other patents pending. No license granted under any patent to use this product other than in accordance with this label. No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE, 1-800-332-3111.
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT, (314)-694-4000.

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF SWALLOWED. HARMFUL IF INHALED. CAUSES MODERATE EYE IRRITATION.
Avoid breathing dust or spray mist. Avoid contact with eyes or clothing.
Remove contaminated clothing and wash clothing before reuse.
Wash thoroughly with soap and water after handling.

INSIDE PAGES (EVEN)

FIRST AID	
IF SWALLOWED	• Call a physician or Poison Control Center for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or physician. • Do not give anything by mouth to an unconscious person. • **Quick treatment is essential to counteract poisoning and should be initiated before signs and symptoms of injury appear.**
IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a Poison Control Center or physician for further treatment advice.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or physician for treatment advice.

• Have the product container or label with you when calling a poison control center or physician, or going for treatment.
• You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information.
• This product is identified as Roundup QuikPRO herbicide, EPA Registration No. 524-535.

DOMESTIC ANIMALS: Keep livestock and pets out of treated areas. Do not graze livestock on treated areas. This product is considered to be relatively nontoxic to digestive and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: Long-sleeved shirt and long pants, protective footwear plus socks, and protective eyewear. Discard clothing and other materials that have been heavily contaminated with this product's concentrate. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations
Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

3.2 Environmental Hazards

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Entry Restrictions: Keep all unprotected persons out of operating areas or vicinity where there may be drift. Keep people and pets off treated areas until spray solution has dried.

INSIDE PAGES (ODD)

4.0 STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container, including rinsate, by application in accordance with label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable Federal, State and local procedures.

CONTAINER HANDLING: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once cleaned, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or Monsanto at 1-800-768-6387. If recycling is not available, puncture and dispose of in a sanitary landfill.

5.0 PRODUCT INFORMATION

Product Description: This product is a postemergent, systemic herbicide with no residual soil activity. It is non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. This product is formulated as a water-soluble granule containing surfactant and no additional surfactant is needed. Apply through most standard sprayers after dissolution and thorough mixing with water according to label instructions.

Time to Symptoms: This product moves through the plant from the point of foliage contact and into the root system. Visible effects on most annual weeds occur within 1 day, and on most perennial weeds in 2 days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a quick yellowing of the foliage which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Mode of Action: One of the active ingredients in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids. A second active rapidly disrupts cell integrity of photosynthetically active tissues in the contacted foliage.

Cultural Considerations: Reduced control may result when application is made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate weed control.

Spray Coverage: Uniform and complete spray coverage will provide best results. Do not spray wood foliage to the point of runoff.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Plants growing from unattached underground rhizomes or root stocks of perennials that have not yet emerged at the time of application will not be affected by the herbicide and will continue to grow.

Annual Maximum Use Rate: The maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as tank mixtures. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds acid equivalent) does not exceed the maximum allowed. For non-crop uses, the combined total of all treatments must not exceed 12.25 pounds of this product (8 pounds of glyphosate acid) per acre per year.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, EXPOSED NON-WOODY ROOTS, OR DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation, since minute quantities of this product can cause severe damage or destruction to desirable plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when weeds are gusty as wind velocity increases, when wind direction is constantly changing or when there are other

INSIDE PAGES (EVEN)

INSIDE PAGES (ODD)

meteorological conditions that affect spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in spatter or fine particles (fines) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or desirable plants, or other unintended consequences. Keep container closed to prevent spills and contamination.

6.0 MIXING

Mix only the amount of solution to be used during a 1-day period. Reduced visual activity but not efficacy will result from the use of leftover solution.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL SEDIMENT IS USED AS A CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS AND DITCHES THAT IS VISIBLY MUDDY OR MURKY.

Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices

where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

6.1 Tank Mixing

This product does not provide residual weed control. This product may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Read and follow all individual product labels, supplemental labeling and Fact Sheets for all products in the tank mixture and observe all precautions and limitations on the label, including application timing restrictions, soil restrictions and use according to the most restrictive precautionary statements for each product in the tank mixture. Always predetermine the compatibility of all tank-mix products together in the carrier by mixing small proportional quantities in advance.

Buyer and all users are responsible for any and all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified on this label. Mixing this product with herbicides or other materials not specified on this label may result in reduced performance.

6.2 Procedure for Preparing Spray Solution

Use the following procedure to mix this product in water alone or when preparing tank mixtures with other labeled products.

1. Place a 20- to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. Add Roundup QuickPRO herbicide using a circular motion while pouring.
4. If second product is a wettable powder, first make a slurry with the water carrier, then add the slurry SLOWLY through the screen into the tank. Continue agitation.
5. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
6. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
7. Continue filling the spray tank with water and add water soluble liquids near the end of the filling process.

When tank mixing this product with other products, maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Use a screen size in nozzle or line strainers no finer than 50-mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

6.3 Mixing for Hand-Held Sprayers

Prepare the desired volume of spray solution by adding the amount of this product as shown in the following table to a clean, empty sprayer. Add the appropriate amount of water and stir or agitate to ensure dissolution of this product. For best results when using backpack sprayers, mix the labeled amount of this product with the specified volume of water in a large container. Fill sprayer with the mixed solution.

Spray Solution

Amount of Roundup QuickPRO herbicide				
Desired Volume	Annuals	Perennials	Brush	Low-Volume Directed
1 Gal	3.2 oz	1.5 oz	1.5 oz	4.0 oz to 8.0 oz
5 Gal	1.6 oz	4.5 oz	4.5 oz	12.0 oz to 1.5 lb
10 Gal	17.0 oz	15.0 oz	15.0 oz	2.5 lb to 5.0 lb

6.4 Colorants or Dyes

Colorants or marking dyes may be added to this product; however, they can reduce product performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's directions. Certain blue dyes are not stable in the spray solution in the presence of this product. A jar test to determine if the desired blue dye is stable is recommended. If stability is a problem consider switching to an alternate color dye.

6.5 Drift Control Additives

Drift control additives may be used with all equipment types. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS.

Avoiding spray drift at the application site is the responsibility of the applicator and grower. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Do not apply this product by air.

Do not apply this product through any type of irrigation system.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to desirable plants or other areas on which treatment was not intended.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

7.1 Ground Broadcast Equipment

Use the labeled rates of this product in 10 to 80 gallons of water per acre as a broadcast spray unless otherwise specified on this label or in separate supplemental labeling or Fact Sheets published for this product. As the density of weeds increases, increase the spray volume within the labeled range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

7.2 Backpack or Hand-Held Equipment

Apply to foliage of vegetation to be controlled on a spray-to-wet basis, do not spray to the point of runoff. For best results, ensure that spray coverage is uniform and complete. Use coarse sprays only.

For control of weeds listed in the **Annual Weeds** section of the **WEEDS CONTROLLED** section, apply 1.7 ounces of this product per 1 gallon of spray solution. See table in **Mixing for Hand-Held Sprayers** section for larger mixing volumes.

For best results, use 1.5 ounces of this product per 1 gallon of spray solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle. See table in **Mixing for Hand-Held Sprayers** section for larger mixing volumes.

For low-volume directed spray applications, use 4.0 to 8.0 ounces of this product per 1 gallon of spray solution for control or partial control of brush weeds. See table in **Mixing for Hand-Held Sprayers** section for larger mixing volumes. Ensure spray coverage is uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of brush and tree seedlings when foliage is thick and dense, or where there are multiple sprouts.

7.3 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount stated in this label when applied by conventional ground equipment. For vehicle-mounted CDA equipment apply 3 to 15 gallons of water per acre.

For hand-held CDA units, apply a solution of 1.5 to 2.0 ounces of this product in one gallon of water at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mile per hour.

CDA equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 SITE AND USE INSTRUCTIONS

Unless otherwise specified, applications may be made to control any weeds listed in the annual, perennial and brush weeds and tree seedlings tables.

8.1 Non-crop Areas and Industrial Sites

Use in non-crop areas only: airports, apartment complexes, commercial sites, Conservation Reserve Program (CRP), ditch banks, driveways, dry ditches, dry canals, fencerows, golf courses, industrial sites, landscape areas, lumberyards, manufacturing sites, municipal sites, natural areas, office complexes, ornamental landscapes, parks parking areas, recreational areas, residential areas, rights-of-way, roadsides, schools, sports complexes storage areas, warehouse areas, and wildlife management areas.

This product is not for use on crops, timber, other plants being grown for sale, other commercial use, or for commercial seed production. This product is not for research purposes.

Weed Control, Trim-and-Edge and Bare Ground

Use this product in non-crop areas. Apply with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation, and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects. This product is not for use on plants grown for sale or other commercial use, or for commercial seed production.

Repeated applications of this product may be used as weeds emerge to maintain bare ground.

When a tank mixture with a generic active ingredient, such as 2,4-D or pendimethalin is described in this label the user is responsible for ensuring that the specific application being made is included on the label of the specific product being used in the tank mixture.

This product may be tank-mixed with the following products. Refer to these products' labels for approved non-crop sites and application rates.

BANVEL	PENDULUM 3.3 EC
BARRICADE 65WG	PENDULUM WDG
CERTAINTY®	RUBISTAR 50 WP
DIMENSION 4 EC	SUREFLAN
ENDURANCE	2,4-D
PENDIMETHALIN	

When applied as a tank mixture for bare ground, Roundup QuickPRO herbicide provides control of the emerged annual weeds and control or partial control of emerged perennial weeds.

Dormant Turfgrass

Use this product to control or suppress many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup. This product is not for use on turf being grown for sale or other

INSIDE PAGES (EVEN)

at least one regular mowing to allow sufficient growth for good interception of the spray.

Do not disturb soil or underground plant parts before treatment. Delay tillage or renovation techniques such as vertical mowing, coring or slicing for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass.

Do not feed or graze treated turfgrass or feed treated thatch to livestock.

8.2 Parks, Recreational and Residential Areas

This product may be used in parks, recreational and residential areas. Apply this product with any application equipment described in this label. Use this product to trim-and-edge around trees, fences, paths, around buildings, sidewalks, and other objects in these areas. This product may be used for spot treatment of unwanted vegetation or to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

All of the instructions in the **Non-Crop Areas and Industrial Sites** section apply to park and recreational areas. This product is not for use around plants being grown for sale or other commercial use.

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre within the range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for rates to control annual and perennial weeds. For difficult to control perennial weeds and where plants are growing under stressed conditions, or where infestations are dense, this product may be used at up to 12.25 pounds per acre for enhanced results.

9.1 Annual Weeds

Use 2.25 to 4.5 pounds per acre of this product as a broadcast spray to control annual weeds. When using rates less than 4.5 pounds per acre, the level of fast-burn symptomology may be reduced.

For spray-to-wet applications, apply 1.2 ounces of this product per 1 gallon of spray solution.

WEED SPECIES

Anoda, spurred	Tarmustardgrass*
Barley*	Little barley*
Barnyardgrass*	London rocket*
Bassia, firehook	Mayweed
Bittercress*	Medusahead*
Black nightshade*	Morningglory (*Ipomoea spp.*)
Bluegrass, annual*	Mustard, blue*
Bluegrass, bulbous*	Mustard, tansy*
Brome, downy*	Mustard, tumble*
Brome, Japanese*	Mustard, wild*
Browntop panicum*	Oats
Bufflegrass*	Pigweed*
Carolina foxtail*	Plains/Tickseed coreopsis*
Carolina geranium	Prickly lettuce*
Castor bean	Purslane, common
Cheatgrass*	Ragweed, common*
Cheeseweed (*Malva parviflora*)	Ragweed, giant
Cheryl*	Red rice
Chickweed*	Russian thistle
Cocklebur*	Rye*
Copperleaf, hophornbeam	Ryegrass*
Corn*	Sandbur, field*
Corn speedwell*	Shattercane*
Crabgrass*	Shepherd's-purse*
Dwarfdandelion*	Sicklepod
Eastern mannagrass*	Signalgrass, broadleaf*
Eclipta*	Smartweed, ladysthumb*
Fall panicum*	Smartweed, Pennsylvania*
Falsedandelion*	Sowthistle, annual
Falseflax, smallseed*	Spanishneedles
Fiddleneck	Speedwell, purslane*
Field pennycress*	Sprangletop*
Filaree	Spurge, annual
Fleabane, annual*	Spurge, prostrate*
Fleabane, hairy (*Conyza bonariensis*)*	Spurge, leafmad*
Fleabane, rough*	Spurry, umbrella*
Florida pusley	Starthistle, yellow
Foxtail*	Stinkgrass*
Goatgrass, jointed*	Sunflower*
Goosegrass	Tansymustard
Grain sorghum (milo)*	Teaweed/Prickly sida
Groundsel, common*	Texas panicum*
Hemp sesbania	Velvetleaf
Henbit	Virginia copperleaf
Horseweed/Marestail (*Conyza canadensis*)	Virginia pepperweed*
Kochia*	Wheat*
Lambsquarters*	Wild oats*
Little barley*	Witchgrass*
London rocket*	Woolly cupgrass*
Mayweed	Yellow nutsedge*
Medusahead*	Yellow rocket
Morningglory (*Ipomoea spp.*)	
Mustard, blue*	
Mustard, tansy*	
Mustard, tumble*	
Mustard, wild*	
Oats	
Pigweed*	
Plains/Tickseed coreopsis*	
Prickly lettuce*	
Purslane, common	
Ragweed, common*	
Ragweed, giant	
Red rice	
Russian thistle	
Rye*	
Ryegrass*	
Sandbur, field*	
Shattercane*	
Shepherd's-purse*	
Sicklepod	
Signalgrass, broadleaf*	
Smartweed, ladysthumb*	
Smartweed, Pennsylvania*	
Sowthistle, annual	
Spanishneedles	
Speedwell, purslane*	
Sprangletop*	
Spurge, annual	
Spurge, prostrate*	
Spurge, leafmad*	
Spurry, umbrella*	
Starthistle, yellow	
Stinkgrass*	
Sunflower*	
Teaweed/Prickly sida	
Texas panicum*	
Velvetleaf	
Virginia copperleaf	
Virginia pepperweed*	
Wheat*	
Wild oats*	
Witchgrass*	
Woolly cupgrass*	
Yellow nutsedge*	
Yellow rocket	

* When using field broadcast equipment (boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled. Applications must be made using 10 to 80 gallons of carrier volume per acre. Use nozzles that ensure thorough coverage of foliage and treat when weeds are in an early growth stage.

INSIDE PAGES (ODD)

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the range.

Use 4.5 to 9.0 pounds per acre of this product as a broadcast spray to control perennial weeds. When using rates less than 9.0 pounds per acre, the level of fast-burn symptomology may be reduced.

For spray-to-wet applications, apply 1.5 ounces of this product per 1 gallon of spray solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

When using hand-held equipment for low-volume directed spot treatments, apply 4.0 to 8.0 ounces of this product per 1 gallon of spray solution.

Allow 7 or more days after application before tillage.

WEED SPECIES

Alfalfa*	Johnsongrass
Alligatorweed*	Kikuyugrass*
Anise (fennel)	Knapweed
Bahiagrass	Lantana
Beachgrass, European (*Ammophila arenaria*)	Lespedeza
Bentgrass*	Milkweed, common
Bermudagrass*	Mullein, common
Bermudagrass, water (Vasegrass)	Napiergrass
Bindweed, field	Nightshade, silverleaf
Bluegrass, Kentucky	Nutsedge, purple, yellow
Bluewed, Texas	Orchardgrass
Bromegrass, smooth	Pampasgrass
Buttonweed, woolly-leaf	Paragrass
Canarygrass, reed	Pepperweed, perennial
Cattail	Phragmites*
Clover, red, white*	Poison hemlock
Cogongrass	Quackgrass
Dallisgrass	Redvine*
Dandelion	Reed, giant
Dock, curly	Ryegrass, perennial
Dogbane, hemp	Sprange, leafy*
Fescue	Thistle, artichoke
Fescue, tall	Thistle, Canada
Geiseman ivy	Timothy
Guineagrass	Torpedograss*
Horsenettle	Trumpetcreeper*
Horseradish	Vaseygrass
Jerusalem artichoke	Velvetgrass
	Wheatgrass, western

* Partial Control

9.3 Brush Weeds and Tree Seedlings

Best results are obtained when brushweeds are treated when they are in the seedling stage of growth. In many situations, retreatment is required on larger plants. Under these conditions, use the higher application rate within the range.

Use 9.0 pounds per acre of this product as a broadcast spray to control brush weeds.

For spray-to-wet applications, apply 1.5 ounces of this product per 1 gallon of spray solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

When using hand-held equipment for low-volume directed spot treatments, apply 4.0 to 8.0 ounces of this product per 1 gallon of spray solution.

Allow 7 or more days after application before tillage.

WEED SPECIES

Alder	Oak, southern red
Ash*	Oak, white*
Beech*	Pepperbrier, Brazilian (Florida holly)*
Birch	Pine
Blackberry	Poison ivy*
Blackgum	Poison oak*
Cherry, bitter, black, pin	Poplar, yellow*
Dogwood*	Redbud, eastern
Elderberry	Rose, multiflora
Elm*	Saltcedar*
Honeysuckle	Sumac, flaxed, poison, smooth, sugarbush, winged*
Locust, black*	Sweetgum
Maple, red	Vine maple*
Maple, sugar	Virginia creeper
Oak, black*	Waxmyrtle, southern*
Oak, northern pin	
Oak, post	
Oak, red	
Oak, scrub*	

* Partial Control

10.0 LIMIT OF WARRANTY AND LIABILITY

Monsanto Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with these Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

To the extent consistent with applicable law, buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company to the extent consistent with applicable law including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather weather conditions which are outside the range considered normal at the application site and for a time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not exactly set forth in the Directions, maintained conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY

INSIDE PAGES (EVEN)

OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Roundup, QuikPRO, DICAMBA, and MonSanto and Vine design are trademarks of Monsanto Technology LLC. All other trademarks are the property of their respective owners.

Product of Brazil, formulated in the U.S. with U.S. Ingredients.

EPA Reg. No. 524-535

In case of an emergency involving this product, Call Collect, day or night. (314) 694-4000

Packed For
MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI, 63167 U.S.A.
©2011
120910

MONSANTO

Roundup QuikPRO Herbicida

Herbicida Roundup QuikPRO es un herbicida profesional no selectivo de acción rápida para utilizar en áreas no cultivadas y sitios industriales.

Instrucciones completas para el uso

EVITE EL CONTACTO DE ESTE HERBICIDA CON FOLLAJE, TALLOS, RAÍCES NO LEÑOSAS EXPUESTAS O PLANTAS Y ÁRBOLES DESEABLES, YA QUE PODRÍA PRODUCIR DAÑOS GRAVES O SU DESTRUCCIÓN.

Registro en la EPA Nº. 524-535 2011-1

Antes de usar este producto, lea la etiqueta en su totalidad.

Úsalo solamente de acuerdo con las instrucciones de la etiqueta.

Antes de comprar o usar el producto, lea "LIMITES EN LA GARANTÍA Y EN LA RESPONSABILIDAD" en la última sección de la etiqueta. Si las condiciones son inaceptables para usted, devuelva el producto inmediatamente sin abrir el recipiente. ESTE ES UN PRODUCTO PARA USARSE TAL Y COMO ESTÁ PREPARADO. MONSANTO NO LO HA DISEÑADO NI LO HA REGISTRADO PARA QUE SEA REFORMULADO NI VUELTO A ENVASAR.

1.0 INGREDIENTES

INGREDIENTES ACTIVO:

*Dicloruro, N-dietilbencenetiólico, en forma de su sal de amonio	73.3%
Diclorurro de dicuat (dibromuro de 6,7-dihidrodipiridio (1,2-a 2',1'-c) pirazinio)	2.9%
OTROS INGREDIENTES	23.8%
	100.0%

*Equivalente a 66.6% de ácido diclorobenzoico.

1.0 libra contiene 0.73 libra de la sal de amonio de diclorobenzoico y 0.03 libra de la sal dibromuro de diquat.

Este producto está protegido por la patente de los Estados Unidos No. 7,008,904. Otras patentes en trámite. No se han otorgado licencias bajo ninguna patente para otros usos del producto que no sean los indicados en esta etiqueta. No se han otorgado licencias bajo ninguna patente que no sea la de estados Unidos.

2.0 TELEFONO IMPORTANTES

PARA INFORMACIÓN SOBRE EL PRODUCTO O AYUDA PARA UTILIZAR ESTE PRODUCTO, LLAME GRATIS AL: 1-800-332-3111.

EN CASO DE QUE SE PRESENTE UNA EMERGENCIA RELACIONADA CON ESTE PRODUCTO, O PARA OBTENER AYUDA MÉDICA, LLAME POR COBRAR A CUALQUIER HORA DEL DÍA O DE LA NOCHE, AL TELÉFONO (314) 694-4000.

INSIDE PAGES (ODD)

3.0 ADVERTENCIAS

3.1 Riesgos para seres humanos y animales domésticos

Manténgase fuera del alcance de los niños.

¡PRECAUCION!

NOCIVO SI SE INGIERE
NOCIVO SI SE PRIVA
PROVOCA IRRITACIÓN OCULAR MODERADA

Evite respirar el polvo o neblinas de aspersión.
Evite el contacto con la piel y con la ropa.
Quítese la ropa contaminada y lávela antes de volver a usarla.
Después de manejar el producto, lávese a fondo con agua y jabón.

PRIMEROS AUXILIOS	
Si es ingerido	• Llame a un médico o a un centro de envenenamientos para que le indique el tratamiento. • Si la persona puede tragar, hágale beber poco a poco un vaso de agua. • No induzca el vómito, a menos que así se lo indique el centro de envenenamientos o el médico. • No administre nada por la boca a una persona que haya perdido el conocimiento. • **Un tratamiento rápido es esencial para contrarrestar la intoxicación** y se debe iniciar antes de que se manifiesten los síntomas y síntomas de daño.
Si es inhalado	• Saque a la persona al aire fresco. • Si la persona no respira, llame al 911 o a una ambulancia, luego administrale respiración artificial, preferentemente boca a boca, si es posible. • Llame a un centro de envenenamientos o a un médico para que le indique el tratamiento posterior.
Si entra en los ojos	• Mantenga abiertos los ojos y enjuáguelos lenta y cuidadosamente con agua durante 15 a 20 minutos. Si está usando lentes de contacto, quítenselos, después de que transcurran 5 minutos, y continúe enjuagando los ojos. • Llame a un centro de envenenamientos o a un médico para que le indique el tratamiento.

- Tenga a mano el envase o la etiqueta del producto cuando llame al centro de envenenamientos o al médico, o cuando vaya para que le indiquen el tratamiento.
- También puede llamar por cobrar al teléfono **(314) 694-4000**, durante el día o la noche, para obtener información sobre el tratamiento médico de urgencia.
- Este producto está certificado como **herbicida Roundup QuikPRO, Nº de Registro de la EPA 524-535.**

ANIMALES DOMÉSTICOS: Mantenga al ganado y a las mascotas fuera de las áreas tratadas. No alimente al ganado en áreas tratadas. Este producto se considera relativamente no tóxico para los perros y otros animales domésticos, sin embargo, la ingestión de este producto o de grandes cantidades de vegetación recientemente tratada puede producir una irritación gastrointestinal temporal (vómitos, diarrea, cólicos, etc.). Si se observan dichas dolencias, déle al animal suficiente cantidad de líquidos para evitar la deshidratación. Si los síntomas continúan por más de 24 horas, llame al veterinario.

Equipo de protección personal (PPE, por sus siglas en inglés)

Los aplicadores y otros tratantes deben usar: Camisa de manga larga y pantalones largos, calzado protector más calcetines, y protección ocular. Deseche la ropa y otros materiales que se han contaminado pesadamente con el concentrado de este producto. Siga las instrucciones del fabricante para la limpieza/mantenimiento del PPE. De no existir dichas instrucciones para elementos lavables, utilice detergente y agua caliente. Mantenga el PPE apartado del resto de la ropa, y lávelo por separado.

Recomendaciones de seguridad para el usuario

Los usuarios deben:

- Lavarse las manos antes de comer, beber, masticar goma, usar tabaco o de usar el inodoro.
- Quitarse de inmediato la ropa si el pesticida pasa a través de la misma. Luego bañarse completamente y ponerse ropa limpia.
- Quitarse el equipo de protección personal (PPE) de inmediato, después de manejar este producto. Lavarse muy bien y ponerse ropa limpia tan pronto como sea posible.

3.2 Riesgos al medio ambiente

Este producto es tóxico para las invertebrados acuáticos. No lo aplique directamente al agua o en áreas donde el agua superficial esté presente o en áreas donde haya mareas altas y bajas, o en donde el nivel medio de mareas altas. No contamine el agua al lavar el equipo o al tirar el agua de lavado.

3.3 Riesgos de orden físico o químico

Para mezclar, almacenar y aplicar la solución de mezcla de este producto, se deben usar solamente recipientes de acero inoxidable, fibra de vidrio, plástico o recipientes de acero recubiertos internamente con fábrica.

INSIDE PAGES (EVEN)

NO MEZCLE, ALMACENE O APLIQUE ESTE PRODUCTO O SUS SOLUCIONES DE ROCIO EN RECIPIENTES O TANQUES RECICLADOS DE ACERO GALVANIZADO O DE ACERO NO RECIBIERTO (EXCEPTO SI ES ACERO INOXIDABLE). Este producto o la solución de rocío reaccionan con el material de dichos recipientes y tanques, lo cual produce hidrógeno, que puede formar una mezcla de gases altamente combustibles. Si esta mezcla de gases entra en contacto con llamas, chispas, el cebo de un soldador, un cigarrillo encendido o cualquier otra fuente de ignición, puede inflamarse o explotar y causar heridas graves a personas.

INSTRUCCIONES PARA EL USO

El uso de este producto de cualquier manera que sea incompatible con las instrucciones dadas en la etiqueta es una violación de las leyes federales. No aplique este producto de alguna manera que entre en contacto con los trabajadores u otras personas, ya sea directamente o por medio del aire. Solamente los aplicadores que usen protección personal pueden estar en el área durante su aplicación. Para verificar requisitos específicos de su hábitat estatal consulte con la agencia responsable de la regulación del uso de pesticidas.

Restricciones de ingreso: Mantenga a todas las personas sin protección alejadas de las áreas de operación o de las áreas próximas en las que pudiera haber deriva. Mantenga a las mascotas y a las mascotas fuera de las áreas tratadas hasta que la solución de rocío se haya secado.

4.0 ALMACENAMIENTO Y DESECHO

Son fundamentales el almacenamiento y la eliminación adecuados de los pesticidas para evitar la exposición de las personas y el medio ambiente como consecuencia de pérdidas y derrames del producto, excedentes o desechos y actos de vandalismo. No permita que este producto contamine el agua, los alimentos de las personas y animales o las semillas con miras al almacenamiento y la eliminación.

ALMACENAMIENTO DEL PESTICIDA: Guarde los pesticidas lejos de los alimentos para personas, los alimentos para mascotas, los alimentos para animales, las semillas, los fertilizantes y los materiales de uso veterinario. Mantenga el envase bien cerrado para evitar derramamientos y contaminación.

ELIMINACION DEL PESTICIDA: Para evitar desechos, use todo el material contenido en este envase, incluyendo los residuos del enjuague, solicitado según las indicaciones de la etiqueta. Si no es posible evitar los desechos, ofrezca el producto restante a una planta de eliminación de desechos o a un programa de eliminación de pesticidas. Estos programas suelen ser manejados por gobiernos estatales o locales o por la industria. Todos los desechos deben seguir los procedimientos federales, estatales y locales aplicables.

MANIPULACION DEL ENVASE: Envase no rellenable. No vuelva a usar este envase para contener materiales que no sean pesticidas o para diluir pesticidas. Después de vaciar y limpiar el envase, en algunos casos se permite su uso para contener temporalmente residuos del enjuague u otros materiales relacionados con los pesticidas. Consulte a su agencia de reglamentación estatal sobre las prácticas permitidas en su estado.

Enjuague tres veces o enjuague por presión el envase (o equivalente) inmediatamente después de una vaciado. Enjuague tres veces de la siguiente forma: Vacíe el contenido restante del envase en el equipo de aplicación o en un tanque de mezcla. Llene el envase con agua hasta un 1/4 de su capacidad. Agite durante 10 segundos. Vierta los residuos del enjuague en el equipo de aplicación o tanque de mezcla o guárdalo para uso o eliminación posterior. Drene durante 10 segundos después de que el flujo comienza a gotear. Repita este procedimiento dos veces más.

Enjuague por presión de la siguiente forma: Vacíe el contenido restante del envase en el equipo de aplicación o en un tanque de mezcla. Sostenga el envase en posición invertida sobre el equipo de aplicación o el tanque de mezcla o recoja los residuos del enjuague para su uso o eliminación posterior. Inserte la boquilla de enjuague por presión en la parte lateral del envase y enjuague a una presión de aproximadamente 40 PSI durante 30 segundos como mínimo. Drene durante 10 segundos después de que el flujo comienza a gotear.

Una vez hecho, algunos envases de pesticida pueden llevarse a un centro de acopio de envases o ser recogidos para su reciclaje. Para encontrar el centro más cercano, póngase en contacto con su distribuidor de productos químicos o con Monsanto en el 1-800-768-6387. Si no es posible reciclar el envase, perfore el envase y deséchelo en un relleno sanitario.

5.0 INFORMACION DEL PRODUCTO

Descripción del producto: Este producto es un herbicida sistémico de aplicación post-emergencia, sin actividad residual en la tierra. No es selectivo y brinda un control de amplio espectro sobre muchas malezas anuales y perennes, malezas leñosas y árboles. Está formulado como un granulado soluble en agua que contiene surfactante y no necesita surfactante adicional. Se puede aplicar con la mayoría de los rociadores estándar. Deslave la tolva diluida y mezcle completamente con agua según las instrucciones de la etiqueta.

Patrón para la aplicación de sistemas: Este producto se desplaza por la planta desde el punto de contacto con el follaje hasta el sistema radicular y dentro de éste. En la mayoría de las malezas anuales los efectos visibles aparecen en 1 día, y en la mayoría de las malezas perennes en 2 días. El clima muy frío o nublado o continuación del tratamiento puede retardar la actividad de este producto y demorar la aparición de síntomas visibles. Los efectos visibles son la rápida aparición de color amarillo en el follaje, que avanza hasta que la vegetación que está por encima de la tierra, quede toda de color marrón, y el deterioro de las partes subterráneas de la planta.

Modo de acción: Uno de los ingredientes activos en este producto inhibe a una enzima que, solo se encuentra en las plantas y en los microorganismos y que es esencial para la síntesis de aminoácidos específicos. Un segundo ingrediente activo rompe rápidamente la integridad de las hojas del follaje con el que entra en contacto, que llevan a cabo la fotosíntesis.

INSIDE PAGES (ODD)

Contraindicaciones culturales: [illegible] Varía resistente si se aplica a malezas anuales o perennes que se han podado, usado para pastoreo cortado, o las que no se les permitió volver a crecer hasta la etapa de tratamiento especificada.

Resistencia a la lluvia: La lluvia torrencial poco después de la aplicación podría lavar el producto del follaje y se necesitaría una nueva aplicación para obtener un control adecuado de las malezas.

Cobertura del rocío: Para obtener mejores resultados, la cobertura del rocío debe ser completa y uniforme. No moje el follaje de la maleza hasta el punto de escurrimiento.

No actividad en el suelo: Las malezas deben haber emergido en el momento de aplicar. En caso de poder ser controladas por este producto, las malezas que germinan de las semillas derivadas de la aplicación no serán controladas. Las plantas que surgen de rizomas subterráneos sin raíces o raíces de perennes que aún no han emergido en el momento de la aplicación no se verán afectadas por el herbicida y continuarán creciendo.

Tasa de aplicación anual máxima: Las cantidades de aplicación máximas permitidas que se aplican a este producto combinado con el uso de cualquier otro herbicida que contenga el ingrediente activo (glifosato, ya sea que se aplique por separado o en otra parte de una mezcla en tanque. Si se aplica más de un producto a base de glifosato al mismo lote dentro del mismo año, debe asegurarse de que el uso total de glifosato (libras de ácido equivalentes) no exceda las proporciones máximas permitidas. En el caso de usos no relacionados con cultivos, el total combinado de todos los tratamientos no debe exceder 12.25 libras de este producto (6 libras de ácido glifosato) por acre por año.

ATENCION

EVITE EL CONTACTO DE ESTE HERBICIDA CON FOLLAJE, RAICES NO LEÑOSAS EXPUESTAS O PLANTAS Y ARBOLES DESEABLES, YA QUE PODRIA PRODUCIR DAÑOS GRAVES O SU DESTRUCCION. EVITE EL ARRASTRE. DEBE PROCEDERSE CON SUMA PRECAUCION AL APLICAR ESTE PRODUCTO PARA EVITAR DAÑAR PLANTAS DESEABLES.

No permita que la solución del herbicida se nebulice, gotee, sea arrastrada o salpique sobre la vegetación deseable. Una cantidad pequeña puede ser suficiente para causar daños graves o destruir las plantas deseables u otras áreas que no se desea tratar. La probabilidad de que ocurran daños por el uso de este producto aumenta cuando hay muchas rafagas de viento, a medida que aumenta la velocidad del viento, cuando la dirección del viento cambia constantemente o cuando existen otras condiciones meteorológicas que favorecen la dispersión del rocío. Cuando se evite aplicando el producto con un rociador, evite la combinación de presiones y bajas de boquilla que puedan dar como resultado salpicaduras o partículas finas (rocío), que tienen muchas posibilidades de que el producto sea arrastrado.

EVITE LA APLICACION A ALTA VELOCIDAD O PRESION EXCESIVAS.

NOTA: El uso de este producto de cualquier forma que no esté indicada en la etiqueta podría producir lesiones a personas y animales o daños a plantas deseables, así como otras consecuencias imprevistas. Mantenga el envase cerrado para evitar derrames y contaminación.

6.0 MEZCLA

Mezcle solo la cantidad de solución que se debe aplicar durante un período de 1 día. El uso de solución remanente reduce la actividad, usualmente hasta la eficacia.

Limpie las cintas del rociador inmediatamente después de su utilización. Herbicidas bien con agua.

NOTA: EL RENDIMIENTO DEL PRODUCTO PODRIA REDUCIRSE SI SE UTILIZA AGUA CON SEDIMENTOS DE TIERRA COMO SUSTANCIA VEHICULAR. NO MEZCLE ESTE PRODUCTO CON AGUA VISIBLEMENTE EMBARRADA O AGUA NO TRANSPARENTE DE ESTANQUES O ACEQUIAS.

Tenga cuidado para evitar el reflujo en la fuente contadora. Utilice aparatos aprobados contra el reflujo en lugares donde lo exijan las normas locales o estatales. Durante la mezcla y aplicación, se puede formar espuma en la solución de rocío. Para prevenir o minimizar la formación de espuma evite el uso de agitadores rociadores cerca de las tuberías de retorno y de pasa en la parte posterior del tanque, y en caso de que sea necesario, utilice un agente antiadhesivo antiespuma o que elimina la espuma.

6.1 Mezcla en tanque

Este producto no proporciona control de malezas resistentes. Este producto puede mezclarse en un tanque con otros herbicidas para lograr el control de malezas resistentes, un espectro de control de malezas más amplio o un modo de acción alternativo. Lea y siga las advertencias incluidas en las etiquetas adicionales, etiquetas suplementarias y fichas de datos de todos los productos de la mezcla de tanque, así como las precauciones y limitaciones incluidas en la etiqueta, reduzca las restricciones en el tanque de aplicación, restricciones del suelo y use según las precauciones más restrictivas para cada producto de la mezcla de tanque. Siempre determine previamente la compatibilidad de todos los productos de la mezcla de tanque. Lo más es la sustancia vehicular, mezclando entre pequeñas cantidades proporcionales.

El combustible y todos los aceites son responsables de cualquier efecto o daño en relación con el uso o manipulación de las mezclas de este producto con herbicidas u otros materiales que no se indican expresamente en esta etiqueta. La mezcla de este producto con herbicidas u otros materiales no indicados en esta etiqueta puede dar como resultado una reducción en el rendimiento del producto.

6.2 Procedimiento para la preparación de la solución de rocío

Utilice el siguiente procedimiento para mezclar este producto solo con agua o cuando prepare mezclas para tanque con otros productos indicados en la etiqueta.

1. Coloque una cantidad de tamaño medio de hielo de 20 a 35 metros el puerto de llenado.
2. A través de la criba, llene el tanque del rociador por la mitad con agua y comience a agitar.
3. Agregue el herbicida Roundup QUAPRO mientras el tanque con un movimiento circular mientras lo vierte.
4. Si el segundo producto es un polvo mojable con agua, proceda haga una pasta con el contenido de frasco, luego agregue LENTAMENTE la pasta a través de la criba dentro del tanque. Continúe agitando.
5. Si se usa una formulación fluida, mezcle previamente una parte fluida con una parte de agua. Agregue la mezcla diluida LENTAMENTE a través de la criba dentro del tanque. Continúe agitando.
6. Si se usa un concentrado emulsionable, mezcle previamente una parte del concentrado emulsionable con dos partes de agua. Agregue LENTAMENTE la mezcla diluida a través de la criba en el tanque. Continúe agitando.
7. Continúe llenando el tanque del rociador con agua y agregue líquidos salientes en agua cerca del final del proceso de llenado.

Al mezclar este producto en un tanque con otros productos, agite bien durante todo el tiempo hasta que el contenido del tanque sea rociado. Si se deja que la mezcla para rociar se asiente, agite bien para que la mezcla vuelva a estar en suspensión antes de continuar con el rociado.

A fin de minimizar la formación de espuma, mantenga las tuberías de retorno lo más cerca del fondo del tanque. La mitad de la rejilla en la boquilla o en los colectores de las mangueras no debería ser de menos de 50 mallas.

INSIDE PAGES (EVEN)

6.4 Colorantes o tinturas

A este producto se le pueden agregar colorantes o tinturas para marcar; sin embargo, pueden reducir su rendimiento, especialmente a bajas concentraciones del producto o a bajas dilusiones. Use colorantes o tinturas según las indicaciones del fabricante. Algunos colorantes pueden ser estables en la solución de rocío en presencia de este producto. Se recomienda una prueba de probeta ("jar test") para determinar si el colorante usado deseado es estable. Si la estabilidad es un problema, considere cambiar a un colorante de color alternativo.

6.5 Aditivos para controlar el arrastre

Se pueden utilizar aditivos que ayuden a controlar el arrastre con todo tipo de equipo. Al utilizar un aditivo para controlar el arrastre, lea y cumpla meticulosamente con las cláusulas preventivas y toda la demás información que aparece en la etiqueta del aditivo.

7.0 EQUIPOS Y TECNICAS PARA LA APLICACION

MANEJO DEL ARRASTRE POR ROCIO

EVITE EL ARRASTRE. DEBE PROCEDERSE CON SUMA PRECAUCIÓN AL APLICAR ESTE PRODUCTO PARA EVITAR DAÑAR PLANTAS DESEABLES.

Evitar el arrastre del rocío en el lugar de aplicación es responsabilidad del aplicador y del agricultor. La interacción de numerosos factores relacionados con los equipos y las condiciones meteorológicas determina el potencial de arrastre del rocío. El aplicador y el agricultor son responsables de tener en cuenta todos estos factores antes de tomar una decisión.

No aplique este producto por aire.

No use ningún sistema de irrigación para aplicar este producto.

No permita que la solución del herbicida se rociadas, gotee, sea arrastrada o salpique sobre la vegetación deseable. Una cantidad pequeña puede ser suficiente para causar daños graves o destruir las plantas deseables u otras áreas que no se desea tratar.

APLIQUE ESTAS SOLUCIONES DE ROCIO UTILIZANDO EQUIPOS FIDEDIGNAMENTE MANTENIDOS Y CALIBRADOS QUE SEAN CAPACES DE ROCIAR EL VOLUMEN DESEADO.

7.1 Equipo de aplicación al voleo terrestre

Use las proporciones de este producto indicadas en la etiqueta en 10 a 40 galones de agua por acre para aplicaciones al voleo a menos que se indique de otra manera en esta etiqueta o en etiquetas suplementarias separadas o en las Fichas de datos publicadas para este producto. A medida que la densidad de las malezas aumenta, el volumen de rocío se debe aumentar tanto en para conseguir una cobertura completa, pero siempre dentro de los límites indicados en la etiqueta. A fin de evitar un rocío muy fino, [illegible] obtener mejores resultados con equipo a nivel del terreno, use boquillas tipo abanico plano. Asegúrese de que las gotas del rocío se distribuyan uniformemente.

7.2 Equipo de mochila o de mano

Aplique al follaje de la vegetación a ser controlada en una aplicación dirigida se rocíe hasta mojar, no rociar al punto de escurrimiento. Para mejores resultados, asegúrese de que la cobertura de rocío sea uniforme y completa. Utilice sólo rociadores gruesos.

Para el control de las malezas enumeradas en la sección del **Malezas anuales** de la sección **MALEZAS CONTROLADAS**, aplique 1.2 onzas de este producto por 1 galón de solución de rocío. Vea la tabla incluida en la sección **Mezcla para rociadores de mano** para volúmenes de mezcla más grandes.

Para obtener mejores resultados utilice 1.5 onzas de este producto por 1 galón de solución de rocío en plantas perennes más difíciles de controlar, como Cynodon dactylon (bermudagrass), Rumex spp. (docal), Convolvulus arvensis (field bindweed), Apocynum cannabinum (hemp dogbane), Aristimoa spp. (milkweed) y Cirsium arvense (Canada thistle). Consulte la tabla en la sección **Mezcla para rociadores de mano** para volúmenes de mezcla más grandes.

En aplicaciones de rocío de gran volumen, use de 4.0 a 6.0 onzas de este producto por 1 galón de solución de rocío para el control o el control parcial de materiales enumeradas en la tabla en la sección **Mezcla para rociadores de mano** para volúmenes de mezcla más grandes. La cobertura del rocío debe ser uniforme y debe entrar en contacto con el 50 por ciento del follaje por lo menos. Para obtener mejores resultados, es importante cubrir la mitad superior de la planta. Para asegurar que el rocío alcance una cobertura adecuada, rocíe ambos lados de las plantas de las materiales las plantas que han crecido de las semillas en los lados cuando el follaje está tupido y espeso, o cuando hay malezas altas.

7.3 Equipo de aplicación por goteo controlado

La tasa de este producto aplicada por aire por el equipo de aplicación por goteo controlado (AGC) no debe ser menor que la cantidad indicada en esta etiqueta cuando se aplica con un equipo al voleo convencional. Cuando se usa el equipo aplicador por goteo controlado montado en un vehículo, use de 3 a 15 galones de agua por acre. Para las unidades de mano del AGC, aplique una solución de 1.5 a 2 litros de este producto en un galón de agua en un caudal de 2 onzas líquidas por minuto y una velocidad de caminada de 0.75 millas por hora.

Los equipos del AGC producen un rocío que es difícil de ver. Se debe tener especial cuidado para evitar que el rocío o el arrastre entre en contacto con el follaje o cualquier otra parte vista de la vegetación deseable, ya que en caso contrario es probable que ésta sea dañada o destruida.

8.0 RECOMENDACIONES SEGUN AREAS Y USO

A menos que se especifique lo contrario, pueden realizarse aplicaciones para controlar las malezas enumeradas en las tablas de malezas anuales, perennes y matorrales y semillas de árboles.

INSIDE PAGES (ODD)

8.1 Áreas no cultivadas y áreas industriales

Utilice solamente en áreas que no se usan para cultivos: perímetros complejos de viviendas, centros comerciales, áreas del Programa de Reservas y Conservación (Conservation Reserve Program, CRP), canales, carriles, las entradas de automóviles, cunetas secas, canales secos, línea de cerca, campos de golf, terrenos industriales, áreas para pistas, decoradas de madera, sitios de telecomunicaciones, sitios municipales, áreas naturales, complejos de oficinas, áreas ornamentales, calzadas, áreas de estacionamiento, áreas recreativas, áreas residenciales, servidumbres de paso, lados de carreteras, estudios, complejos deportivos, líneas de dispositivos, líneas de almacenamiento y líneas de transmisión de vida silvestre.

Este producto no debe usarse en cultivos agrícolas, cultivos de árboles para madera, otras plantas que se cultivan para vender, otros usos comerciales o para la producción comercial de semillas. Este producto no debe usarse para investigación.

Control de malezas, recortes y bordes y suelo limpio de malezas

Utilice este producto en áreas no cultivadas. Aplique con rociadores de los equipos descritos en esta etiqueta.

Puede usarse para el recortado de bordes alrededor de árboles en áreas no cultivadas, para tratamiento localizado de vegetación no deseable y para eliminar las malezas no deseables que crecen en cuadros de árboles establecidos y plantaciones ornamentales. Este producto puede usarse antes de plantar un área con plantas ornamentales, flores, césped (tepes o semillas), o antes de colocar asfalto o de comenzar un proyecto de construcción. Este producto no se debe usar en plantas que se cultivan para la venta o otro uso comercial, ni para la producción de semillas comerciales. Puede hacerse aplicaciones repetidas de este producto a medida que emergen las malezas para mantener el suelo limpio de malezas.

Cuando en esta etiqueta se recomienda una mezcla de tanque con un ingrediente activo genérico como 2,4-D, o herbicida, el usuario asume la responsabilidad de asegurarse de que la solución específica que está preparando está incluida en la etiqueta del producto que está utilizando en la mezcla de tanque. Este producto puede ser mezclado en el tanque con los productos siguientes. Refiérase a las etiquetas de estos productos para los usos aprobados de las áreas no cultivadas y las proporciones de uso.

BANVEL	PENDIMETHALIN 3.3 EC
BARRICADE 65WG	PENDULUM WDG
CERTAINTY®	RONSTAR 50 WP
DIMENSION 4 EC	SURFLAN
ENDURANCE	2,4-D
FENTHIMETHALIN	

Cuando es aplicado como mezcla para el suelo limpio de malezas, el herbicida Roundup QuickPRO proporciona el control de las malezas anuales emergentes y el control o control parcial de las malezas perennes emergentes.

Césped durmiente

Use este producto para controlar o suprimir muchas malezas anuales de invierno y tallo fresco (Festuca arundinacea) para el alivio efectivo de césped de bermudagrass y bahiagrass durmientes. Trate solamente cuando el césped está durmiente y antes de levantarse en la primavera. Este producto no se debe usar en césped que se cultiva para la venta o otro uso comercial, como tepes, ni para la producción de semillas comerciales, ni con fines de investigación.

Aplique entre 5 y 16 onzas de este producto por acre. Aplique las proporciones indicadas en la etiqueta en 10 a 40 galones de agua por acre. Utilícelo sólo en áreas donde bermudagrass o bahiagrass sean cobertura de terreno deseables y donde pueda tolerar algún daño temporal o la decoloración.

Los tratamientos que superen las 9 onzas por acre pueden dar como resultado el daño o la demora para verdear en áreas muy conservadas, como campos de golf o césped.

Renovación de césped (Excepto en plantaciones comerciales de tepes)

Este producto controla la maleza de la vegetación existente antes de la renovación de áreas de césped. Este producto no se debe usar en césped que se cultiva para la venta u otro uso comercial como tepes, ni para la producción de semillas comerciales, ni con fines de investigación. Para lograr el máximo control de la vegetación existente, pertinaz, la siembra o la cosechada con césped de un terreno para determinar si es posible algún retrono a partir de partes subterráneas de plantas que se salvaron. En las áreas donde sea necesario repetir los tratamientos, debe haber suficiente crecimiento antes de la aplicación. No use este producto para la renovación de céspedes de las especies Cynodon dactylon (bermudagrass) y Zoysiatenuifolia (Korean grass). En los lugares donde la vegetación existente crece bajo un manejo de corte periódico, el uso este producto después de remitir al menos una siega habitual de modo que crezca la suficiente para que el rocío actúe con eficacia.

No pertube la tierra o las partes subterráneas de las plantas antes de realizar el tratamiento. Las técnicas de labranza o renovación como segado vertical, ahuecamiento o rotarado deberían retrasarse por 7 días después de la aplicación para que no produzca el correcto traslado del producto a las partes subterráneas de la planta.

Puedan plantarse los céspedes deseados siguiendo los procedimientos anteriormente mencionados.

Puede utilizarse un equipo de mano para el tratamiento localizado de vegetación no deseada que crezca en el césped existente.

No pastoree ni alimente al ganado con el césped tratado, ni lo alimente con paja de heno tratado.

8.2 Parques, áreas recreativas y residenciales

Este producto puede usarse en parques, áreas recreativas y residenciales. Aplique este producto con cualquiera de los equipos descritos en esta etiqueta. Use este producto para el recortado de bordes alrededor de árboles, vallas, caminos, alrededor de edificios, aceras y otros objetos en estas áreas. Este producto puede usarse para tratamiento localizado de vegetación no deseable o para eliminar las malezas no deseables que crecen en lechos de arbustos establecidos y plantaciones ornamentales. Este producto puede usarse antes de plantar un área con plantas ornamentales, flores, césped (tepes o semillas), o antes de colocar asfalto o de comenzar un proyecto de construcción.

Todas las instrucciones de la sección Áreas no cultivadas y Áreas industriales son válidas para los parques y áreas recreativas. Este producto no se debe usar alrededor de plantas que se cultivan para la venta ni para otro uso comercial.

INSIDE PAGES (EVEN)

9.0 TIPOS DE MALEZAS CONTROLADAS

Use siempre la proporción más alta de este producto por acre, dentro de las proporciones, cuando las malezas son densas o cuando crecen en un área no tratada (no cultivada). Puede haber una disminución de los resultados cuando se dañan malezas cubiertas con mucho polvo. Para las malezas que han sido segadas, pastadas o cortadas, permita que vuelvan a crecer antes del tratamiento. Referirse a las secciones siguientes para las proporciones para el control de malezas anuales y perennes. Para las malezas perennes difíciles de controlar y donde las plantas están creciendo bajo condiciones agrestes, o donde hay infestaciones intensas, este producto se puede usar en una concentración de hasta a 12.25 libras por acre para obtener mejores resultados.

9.1 Malezas anuales

Use de 2.25 a 4.5 libras por acre de este producto para aplicación al vuelo para controlar malezas anuales. Cuando se utilizan proporciones menores de 4.5 libras por acre, el nivel de sintomatología de desarrollo rápido puede reducirse.

Para aplicaciones de inicio para mejor, agregue 1.7 onzas de este producto por 1 galón de solución de inicio.

ESPECIES DE MALEZAS

Acedo, sparrel	Lambsquarters*
Barley*	Little barley*
Bermudagrass*	London rocket*
Bassia, fireweed	Maywead
Bitterweed*	Medusahead*
Black nightshade*	Morningglory (Ipomoea spp.)
Bluegrass, annual*	Mustard, blue*
Bluegrass, bulbous*	Mustard, tansy*
Brome, downy*	Mustard, tumble*
Brome, Japanese*	Mustard, wild*
Brownstop panicum*	Oats
Buttercup*	Ragweed*
Carolina foxtail*	Plains/Tall cheat (cheatgrass)*
Cardaria pubescens	Prickly lettuce*
Carpet bean	Purslane, common
Chealgrass	Ragweed, common*
Cheatweed (Malva parviflora)	Ragweed, giant
Chicory*	Redbrome
Crabweed*	Russian thistle
Cocklebur*	Rye*
Coolpetbur, Inapertinbur	Ryegrass*
Corn*	Sandbur, field*
Corn speedwell*	Shattercane*
Crabgrass*	Shepherd's-purse*
Dwarf dandelion*	Sicklepod
Eastern mannagrass*	Signalgrass, broadleaf*
Eclipta*	Smartweed, ladysthumb*
Fall panicum*	Smartweed, Pennsylvania*
Falsedandelion*	Sowthistle, annual
Falsaflax, smallseed*	Spanishneedles
Fiddleneck	Standard, purslane*
Field pennycress*	Sprangletop*
Filaree	Spurge, annual
Fleabane, annual*	Spurge, prostrate*
Fleabane, hairy (Conyza bonariensis)*	Spurge, spotted*
Fleabane, rough*	Spurry, umbrella*
Florida pusley	Starthistle, yellow
Foxtail*	Stinkgrass*
Goatgrass, jointed*	Sunflower*
Goosegrass	Tansymustard (Hedge)
Grain sorghum (Hedge)*	Tansy ragwort*
Groundsel, common*	Velvetleaf
Hemp sesbania	Virginia copperleaf
Henbit	Virginia pepperweed*
Horseweed, Marestail (Conyza canadensis)	Wheat*
Italiangrass*	Wild oats*
Johnsongrass, seedling	Wildgrass*
Junglerice	Woolly cupgrass*
Knotweed	Yellow rocket
Kochia	

*Cuando se realiza de aplicación al vuelo a nivel del terreno (boquillas de aguilón con hoja flat fan blanco plano), estas especies serán controladas o controladas parcialmente. Las aplicaciones deben hacerse un mínimo de 10 a 60 galones de volumen por acre. Use boquillas que garanticen una cobertura completa del follaje y haga el tratamiento cuando las malezas estén en su etapa temprana de crecimiento.

9.2 Malezas perennes

Los mejores resultados se obtienen cuando se trata a las malezas perennes después de que alcanzan la etapa reproductiva (inicio de formación de semillas en hierbas y formación de yemas en malezas de hoja ancha). En las plantas sin flores, los mejores resultados se obtienen cuando las plantas alcanzan la "roseta". En muchas situaciones es necesario realizar tratamientos antes de esas etapas. En esas condiciones, use la dosis de aplicación más alta dentro del rango.

Use de 4.5 a 9.0 libras por acre de este producto para aplicación al vuelo para controlar malezas perennes. Cuando se utilizan proporciones menores de 9.0 libras por acre, el nivel de sintomatología de desarrollo rápido puede reducirse. Para las aplicaciones de inicio para mejor, agregue 1.5 onzas de este producto por 1 galón de solución de inicio. Asegúrese de lograr una cobertura completa cuando efectúe tratamientos de inicio para mejor mediante un equipo de mano.

Cuando utilice un equipo de mano para tratamientos localizados dirigidos de bajo-volumen, agregue de 4.0 a 8.0 onzas de este producto por 1 galón de solución de inicio.

Espere 7 días o más después de la aplicación antes de labrar.

ESPECIES DE MALEZAS

Alfalfa*	Johnsongrass
Alligatorweed*	Knotgrass*
Aster (fetwort)	Knotweed
Bahiagrass	Lantana
Beachgrass, Dunegrass (Ammophila arenaria)	Leafy spurge
Bentgrass*	Milkweed, common
Bermudagrass*	Musky wintertem
Bermudagrass, water (Dallisgrass)	Mullein, common
Brookweed, Tutti	Nutsedgrass
Bluegrass, Kentucky	Nightshade, silverleaf
Bluewead, Texas	Nutsedge, purple, yellow
Brainrgrass, smooth	Orchardgrass
Burdoque, woolly-leaf	Pennsylvania
Canarygrass, reed	Paragrass
Cattail	Peppermwed, perennial
Clover, red, white*	Pineapplegrass*
Cogongrass	Pokeweed
Dallisgrass	Quackgrass
Dandelion	Redwire*
Dock, curly	Reed, giant
Dogbane, hemp	Ryegrass, perennial
Fescue	Sourgrass, tall*
Fescue, tall	Thistle, artichoke
German ivy	Thistle, Canada
Guineagrass	Timothy
Horsenettle	Torpedograss*
Horseradish	Trumpetcreeper*
Kikuyu	Vaseygrass
Knotgrass	Velvetgrass
Jerusalem artichoke	Wheatgrass, western

*Control Parcial

INSIDE PAGES (ODD)

9.3 Matorrales y plántulas de árboles

Los mejores resultados se obtienen cuando los matorrales se tratan en la etapa de plántula. En muchas ocasiones es necesario volver a tratar las plantas más grandes. Bajo estas condiciones, use las proporciones de aplicación más alta dentro del rango.

Para controlar matorrales use 9.0 libras por acre de este producto en forma de aplicación al vuelo.

Para aplicaciones de inicio para mejor, agregue 1.5 onzas de este producto por 1 galón de solución de inicio. Asegúrese de lograr una cobertura completa cuando efectúe tratamientos de inicio para mejor mediante un equipo de mano.

Cuando utilice un equipo de mano para tratamientos localizados dirigidos de bajo-volumen, agregue de 4.0 a 8.0 onzas de este producto por 1 galón de solución de inicio.

Espere 7 días o más después de la aplicación antes de labrar.

ESPECIES DE MALEZAS

Alder	Oak, southern red
Ash*	Oak, white*
Beech*	Persimmon, Brazilian (Florida holly)*
Birch	Pine
Blackberry	Poison ivy*
Blackgum	Poison oak*
Cherry, bitter	Poplar, yellow*
Black, pin	Redbud, eastern
Dogwood*	Rose, multiflora
Elsberry	Saltcedar*
Elm*	Sumac, flameleaf, poison, smooth, sugarbush, winged*
Hickory, shellbark	Sweetgum
Locust, black*	Vine maple*
Maple, red	Virginia creeper
Maple, sugar	Wayfaring tree, southern*
Oak, black*	
Oak, northern pin	
Oak, post	
Oak, red	
Oak, scrub*	

*Control parcial

10.0 LIMITES EN LA GARANTIA Y EN LA RESPONSABILIDAD

Monsanto Company garantiza que este producto concuerda con la descripción química de la etiqueta y es razonablemente adecuado para los propósitos descritos en el folleto "Indicaciones Completas para el Uso" ("Instrucciones") cuando se usa de acuerdo con dichas Instrucciones y las condiciones que allí se detallan.

NO SE HACE NINGUNA OTRA GARANTIA EXPRESA O IMPLICITA ACERCA DE LA IDONEIDAD PARA UN USO IMPARTICULAR O COMERCIABILIDAD.

Esta garantía está sujeta tanto en a las condiciones y limitaciones que aquí se indican.

Hasta el grado en que sea compatible con la legislación pertinente, el comprador y todos los usuarios deberán notificar inmediatamente a esta Compañía acerca de cualquier reclamo que se base en un contrato, negligencia, estricta responsabilidad u otros defectos extracontractuales.

Hasta el grado en que sea compatible con la legislación pertinente, el comprador y todos los usuarios son responsables por todas las pérdidas o daños que resultasen por el uso o manipulación en condiciones que estén más allá del control de esta Compañía, incluyendo pero no limitándose a incompatibilidad con productos que no sean los señalados en las Instrucciones, aplicación o contacto con vegetación susceptible, condiciones climáticas inusuales, condiciones de clima que estén fuera de los límites que se mencionan o normales en el lugar de la aplicación y para el período de tiempo en el cual se aplica, así como condiciones de clima que estén fuera de los límites indicados en las Instrucciones, aplicaciones que no estén específicamente acompañadas en las Instrucciones, condiciones de humedad que estén fuera de los límites establecidos en las Instrucciones, o la presencia de productos en la tierra o sobre ella, en los cultivos o en la vegetación que no sea natural, diferentes a los indicados en las Instrucciones.

Monsanto Company no garantiza ninguna de los productos reformulados o temporizados de este producto, excepto de acuerdo a los requisitos de la administración de esta Compañía y con el permiso escrito expreso de esta Compañía.

HASTA EL GRADO EN QUE SEA COMPATIBLE CON LA LEGISLACIÓN PERTINENTE, LA ÚNICA Y EXCLUSIVA COMPENSACIÓN AL USUARIO O COMPRADOR Y EL LÍMITE DE RESPONSABILIDAD DE ESTA COMPAÑÍA O DE CUALQUIER OTRO VENDEDOR POR CUALQUIER PÉRDIDA O POR TODAS LAS PÉRDIDAS, PERJUICIOS O DAÑOS QUE RESULTASEN DEL USO O MANEJO DE ESTE PRODUCTO (INCLUYENDO RECLAMOS QUE SE BASEN EN UN CONTRATO, NEGLIGENCIA, ESTRICTA RESPONSABILIDAD Y U OTROS DERECHOS EXTRACONTRACTUALES.) SERÁ EL PRECIO PAGADO POR EL USUARIO O EL COMPRADOR POR LA CANTIDAD INVOLUCRADA DE ESTE PRODUCTO, O A ELECCIÓN DE ESTA COMPAÑÍA O DE OTRO VENDEDOR, EL REEMPLAZO DE DICHA CANTIDAD O SI O SE OBRUVO MEDIANTE COMPRA SE REEMPLAZARÁ DICHA CANTIDAD DEL PRODUCTO.

HASTA EL GRADO EN QUE SEA COMPATIBLE CON LA LEGISLACIÓN PERTINENTE, EN NINGÚN CASO ESTA COMPAÑÍA U OTRO VENDEDOR SERÁN RESPONSABLES POR DAÑOS INCIDENTALES, CONSECUENTES O ESPECIALES.

En el momento de abrir y usar el producto, se asume que el comprador y todos los usuarios han aceptado las condiciones de los LIMITES EN LA GARANTIA Y EN LA RESPONSABILIDAD que no pueden variar por medio de origen acuerdo verbal o escrito. Si las condiciones son inaceptables, devuelva el producto inmediatamente sin abrir el recipiente.

Certa-ety, Roundup QUICKPRO y Monsanto y el símbolo de la vid son marcas registradas de Monsanto Technology LLC. Todas las otras marcas registradas son propiedad de sus respectivos dueños.

Registro en la EPA Nº 524-535

En caso de que se presente una emergencia relacionada con este producto, llame por cobrar al número en horas del día o de la noche, al teléfono (314) 694-4000.

Producto de Brasil, formulado en EE.UU. con ingredientes de los EE.UU.

Envíe todo por:
MONSANTO COMPANY
800 N. LINDBERGH BLVD.
ST. LOUIS, MISSOURI 63167, EE.UU.
©2009
120810

MONSANTO

BASE COPY

FIRST AID

If SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. • Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-441-3637 for information.
If INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-nose if possible. • Call a poison control center or doctor for further treatment advice.
If in EYES	• Hold eyelid open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

• Have the product container or label with you when calling a poison control center or doctor or going for treatment.

• For emergency assistance call 1-800-332-4009 collect or contact your local emergency medical treatment facility.

• For information contact the Monsanto 1-800-Monsanto (800-766-7266) or EA Registration Inc. 1-800-345-3275.

Dry Salt Water Recommendation

Dry Salt Water

• Add 1 lb of salt per gallon of water. Stir and allow solution to settle. Decant the liquid. Remove 50% of solution by pouring off or pumping. Stir remaining solution and pour into a drain.

• Remove 75% of solution by pouring off or pumping. Stir remaining solution and pour into a drain.

• Remove 75% of solution by pouring off or pumping. Stir remaining solution and pour into a drain.

Environmental Hazards

This product is toxic to fish. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. For sale only to persons who will use the product for the purpose and in the manner described on the label. Do not use this product in greenhouses or other enclosed areas. Do not use this product in a manner that may result in injury to humans, domestic animals, or other non-target organisms. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not apply this product through any irrigation system. Do not apply this product through any irrigation system. Do not apply this product through any irrigation system.

Storage and Disposal

Pesticide storage: Store in original container in a secure, dry place to prevent leakage or spillage and to prevent contamination of other products, feed, or foodstuffs. Do not store near feed or foodstuffs. PESTICIDE STORAGE: Store in original container in a secure, dry place to prevent leakage or spillage and to prevent contamination of other products, feed, or foodstuffs. Do not store near feed or foodstuffs. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse (or equivalent) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

See back panel for complete STORAGE and DISPOSAL language.

(See back panel for complete PRECAUTIONS, PERSONAL PROTECTIVE EQUIPMENT and COMPLETE DIRECTIONS FOR USE)

MONSANTO

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION! [illegible]

[illegible]

[illegible]

[