

Feb 23, 2024

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

DOC ID# 592206

Listed by the Organic Materials Review Institute (OMRI) for use in organic production

FOR ORGANIC PRODUCTION

The active ingredient in this product is exempt from the requirement for a tolerance when used as a molluscicide in or on all food commodities. May be used up to and including day of harvest.

ACTIVE INGREDIENT	By Wt.
Iron phosphate	1.0%
OTHER INGREDIENTS	99.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the hotline number 1-800-255-3924

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE) Requirements:

Applicators and other handlers must wear: long-sleeved shirts and long pants; and shoes plus socks. Follow manufacturer's instructions for maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

GENERAL INFORMATION

- This product contains a unique combination of iron phosphate, which is a plant nutrient that occurs in soils, with slug and snail bait additives. The bait which is not ingested by snails and slugs will degrade and become a part of the soil.
- The bait is ingested by slugs and snails when they travel from their hiding places to plants. Ingestion, even in small amounts, will cause them to cease feeding. This physiological effect of the bait gives immediate protection to the plants even though the slugs and snails may remain in the area. After eating the bait, the slugs and snails cease feeding, become less mobile and begin to die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease in plant damage.
- This product will control snails and slugs in greenhouses, outdoor ornamentals, vegetable gardens, fruits, berries, citrus and crop plants. The bait can be scattered on the lawn or on the soil around any vegetable plants, flowers or fruit trees or bushes where snail or slug control is needed.
- This product is effective against a wide variety of slugs and snails including (but are not limited to): *Deroceras reticulatum* (Field slug), *Deroceras laeve* (Smooth slug), *Arion subfuscus* (Dusky slug), *Arion circumscriptus* (Gray garden slug), *Arion hortensis* (Black field slug), *Arion rufus* (Large red slug), *Arion ater* (Large black slug), *Limax flavus* (Spotted garden slug), *Limax tenellus* (Slender slug), *Ariolimax columbianus* (Banana slug), *Helix* spp., *Helicella* spp., and *Cepaea* spp.
- Remains effective after rain or sprinkling.
- Can be used around domestic animals and wildlife.

Sluggo®

SLUG AND SNAIL BAIT



NET WEIGHT 50 lbs

EPA Registration No.67702-3-70051
EPA Establishment No.48498-CA-2
Lot Number:

Manufactured for
Certis USA LLC
9145 Guilford Road, Suite 175
Columbia, MD 21046

CERTIS



Sold under a license of
W. Neudorff GmbH KG
An der Mühle 3
31860 Emmerthal – Germany

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a manner that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency representative responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and the restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

There is a restricted-entry interval of zero hours for this product.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

There is a restricted-entry interval of zero hours for this product.

COMMERCIAL AGRICULTURE

HOW TO APPLY: Scatter the slug and snail bait granules on the soil around or near the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply the higher rates if the infestation is severe or if the area is heavily watered or after long periods of heavy rain. Reapply as the bait is consumed or at least every two weeks.

Do not place in piles. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. See specific directions for different plant types and for inside greenhouses.

WHEN TO APPLY: Apply the bait in the evening, as slugs and snails travel and feed mostly by night or early morning.

WHERE TO APPLY: Treat all likely areas of infestation, especially around the perimeter of garden plots because these pests travel into plant areas from daytime refuge sites. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery and squash, and in weeds or ditches around field margins. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day.

Vegetables

Use the bait to protect any vegetables from slug and snail damage, including (but not limited to): asparagus, beans, beets, blackeyed peas, broccoli, Brussels sprouts, cabbage, cantaloupe, carrots, cauliflower, corn, cucumbers, eggplants, garlic, lettuce, onions, peas, peppers, potatoes, radishes, rutabagas, spinach, squash, Swiss chard, tomatoes and turnips. Scatter the bait around the perimeter of the vegetable plantings at the rate of 20-44 lbs. per acre (0.5-1 lb. per 1,000 square feet) to provide a protective "barrier" for slugs and snails entering the vegetable plantings. If slugs or snails are inside the rows, then scatter the bait on the soil around the plants and between the rows.

Fruits Including Citrus

Use the bait to protect fruits from slugs and snails, including (but not limited to): apples, avocados, apricots, cherries, grapes, melons, peaches, plums, nectarines, citrus, pears. For seedlings, spread the bait around the base of the stem. Apply at 0.15 oz., or 1 level teaspoon, per square yard, in a 6-inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the trees to intercept slugs and snails traveling to the trunk. Apply the bait at 20-44 lbs. per acre (0.5-1 lb. per 1,000 square feet) in orchards, using standard fertilizer granular spreaders. Use the higher rates for heavy infestations.

Berries

Use the bait to protect berries from slugs and snails, including (but not limited to): strawberries, blackberries, blueberries, boysenberries, loganberries, raspberries. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of 20-44 lbs. per acre (0.5-1 lb. per 1,000 square feet) and scatter by hand or with granular spreaders. If slugs and snails are already in the plots, then carefully spread bait between the furrows near the base of the plants. For small plots, treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

Field Crops

Use the bait to protect field and seed crops from slugs and snails, including: beans, field corn, sweet corn, soybeans, sugarbeets, sugar cane, asparagus, beets, broccoli, Brussels sprouts, cabbage, carrots, cauliflower, cucumbers, lettuce, onions, peas, peppers, potatoes, radishes, strawberries, tomatoes, and turnips. At the seedling and later stages, apply the bait between the rows and around the perimeter of the field. Scatter pellets at a rate of 20-44 lbs. per acre (0.5-1 lb. per 1,000 square feet). Use the higher dosage rate for heavy infestations.

Artichokes

Use the bait to protect artichokes from slugs and snails. At the seedling and later stages, apply the bait within the rows and around the perimeter of the field. Scatter pellets at a rate of 10-44 lbs. per acre (0.23-1 lb. per 1,000 square feet). Use the higher dosage rate for heavier infestations.

Outdoor Ornamentals

Scatter bait in a 6-inch circular band around the base of the plants to be protected at 0.15 oz., or 1 level teaspoon, per square yard. If plants are next to a grassy area, spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants will encounter the bait before reaching the plant. In these situations, scatter the bait around the perimeter of the plot at 1 lb. per 1,000 square feet to intercept snails and slugs traveling to the plot.

Greenhouses

Where slugs or snails are a problem in greenhouses, scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply about ½ teaspoon per 9-inch pot.

Outdoor Container-Grown Nursery Plants

Where slugs or snails are a problem in outdoor nurseries, scatter the bait in the plant containers at the rate of 1 tablespoon per container of plants being damaged, or scatter around the soil near the containers at the rate of 1 teaspoon per square yard (24-44 lbs. per acre).

Commercial Turf Sod, Golf Courses and Grass Buffers around Gardens, Crop Areas and Ornamentals

Use the bait on turf and sod for resale, golf courses and on lawns as a buffer zone around gardens and agricultural crop areas. When slugs or snails are detected, scatter the bait at a rate of 0.54 to 1 lb. per 1000 square feet (24-44 lbs. per acre). Use the higher rate for heavy infestations. Scatter by hand or with a granular spreader where the slugs or snails are observed.

Grass Grown for Seed Production, Wheat

Use the bait to protect grass grown for seed production and wheat. When slugs or snails are detected, scatter the bait at a rate of approximately 0.23 to 1 lb. per 1,000 square feet (10-44 lbs. per acre). Use the higher rate for heavy infestations. Scatter by hand or with a granular spreader where the slugs or snails are observed.

Surrounding Agricultural Areas Suffering Infestation

Use the bait to protect non-crop areas, including: fallow land, barrier strips and buffer zones around agricultural crop areas. Apply at a rate of 20-44 lbs./acre (0.5-1 lb. per 1,000 square feet). Use the higher rate for heavy infestations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty metal drum, plastic bag, box or plastic tote into application equipment. Then dispose of empty metal drum, plastic bag, box or plastic tote in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

To the extent consistent with applicable law, the seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated on this label only when used in accordance with the directions for use. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. To the extent consistent with applicable law, seller makes no other warranties, either expressed or implied.



ESL 20140908
Ver. 20240206

152205