

FIFRA 2(ee) Recommendation For Management of Conifer Blights in Field Production, Nursery and Outdoor Landscapes

Torque™ Fungicide
EPA Reg. No. 1001-87

This recommendation is made as permitted under FIFRA Section 2(ee)
and has not been submitted to or accepted by the USEPA

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

PLANTS	DISEASE	APPLICATION
Conifers • Juniper[s] • Pine[s]	Phomopsis Blight <i>Phomopsis juniperovora</i>	For Conifer Blights, apply at a rate of 6-10 fl. oz./100 gallons as a full coverage spray to the point of drip. Apply as needed, beginning when conditions are favorable for disease development.
	Tip Blight <i>Sirococcus strobilinus</i>	
	Diplodia Tip Blight <i>Sphaeropsis sapinea</i>	For Diplodia Tip Blight and Tip Blight, apply every 14-21 days during the growing season, just prior to expected infection window, repeat as needed based on weather and disease pressure. Application should be started at the initiation of new growth. For Juniper Phomopsis Blight, apply every 14-21 days during the growing season, just prior to expected infection window, repeat as needed based on weather and disease pressure. Make first application as soon as junipers start to grow. Can be used as a tank mix or rotation with 3336, Spectro, or Chlorothalonil.

ACCEPTED

March 18, 2026

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc id: 605042

Classified for
“RESTRICTED USE”
in New York State
under 6NYCRR Part 326