

March 23, 2026

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

FIFRA 2(ee) Recommendation For Disease Management

In Field Production, Nursery and Outdoor Landscapes

Classified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

Doc id: 605138

Torque™ Fungicide EPA Reg. No. 1001-87

This recommendation is made as permitted under FIFRA Section 2(ee)
and has not been submitted to or accepted by the USEPA

ORNAMENTAL DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, just prior to expected infection window, repeat as needed based on weather and disease pressure.	Apply every 14-21 days beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply every 14-21 days just prior to expected infection window, repeat as needed based on weather and disease pressure.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthrachnose Leaf Spot Powdery Mildew Rust Scab	Apply every 14-21 days beginning with spring bud break or just prior to expected infection window, repeat as needed based on weather and disease pressure.	
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs Ground Covers and Vines	Anthrachnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust Southern Blight	Apply every 14-21 days beginning with spring bud break or just prior to expected infection window, repeat as needed based on weather and disease pressure. Petal Blight – Apply 2-3 times per week into the flowers as they open and develop color.	