

Japanese Knotweed

Identification, Biology and Management

By Drew Zwart, PhD, Plant Physiologist

Japanese knotweed, *Fallopia japonica* (synonyms = *Reynoutria japonica* = *Polygonum cuspidatum*) is a major invasive plant problem throughout North America, and is listed by the World Conservation Union as one of the “world’s worst invasive species.” Native to eastern Asia, the plant is aggressively invasive and causes many environmental and landscape concerns.

Identification

Japanese knotweed is an herbaceous or semi-woody, hollow-stem perennial with deciduous foliage. Identifying characteristics include heart- or spade-shaped leaves with flat bases, prolific small flowers that are white to light green and develop in leaf axils and near the ends of branches, and a brown bamboo-like appearance in winter. Knotweed has raised nodes on the stems, similar to bamboo, and newly developing leaves and stems often have a red to purple coloration.

Concerns

As an aggressive invader, knotweed displaces native plants through competition for light and nutrients (Figure 1), and possible allelopathy (chemical toxicity

Figure 1: Japanese knotweed invading a lawn



Figure 2: Japanese knotweed invading a lake shore



to other plants). In addition to negative effects on native plants, knotweed grows prolifically in watersheds and can alter hydrology, negatively impact habitat of aquatic organisms, and cause many related ecological problems throughout an invaded habitat. Knotweed invasions often spread downstream as

water carries small pieces of plants which start new infestations wherever they are deposited (Figure 2). Knotweed also produces deep and aggressive roots and rhizomes (root-like storage organs). These underground parts can damage building foundations, walls, driveways, sidewalks, and any other built structures. Shoots of knotweed have even been documented to sprout directly through concrete and asphalt.

Management

Physical removal of Japanese knotweed, mowing, or other mechanical means of control are very rarely effective because of its deep root system and ability to sprout from tiny vegetative fragments. In some cases, physical removal actually results in a larger plant or patch of plants the following year. Even complete removal of 2-3 meters of soil (to remove roots and rhizomes) has proven ineffective in many cases. Composting knotweed is not an effective eradication method and is more likely to spread the plant.

Successful management requires multiple applications of a systemic herbicide that moves into and kills root systems, preventing re-sprouting. Even with this approach, several years of treatment are usually required to eradicate a population. Foliar applied systemic materials can be effective if employed correctly. Once the plant is dead, the debris can be removed without fear of spread. For heavily infested sites or older well-established populations (Figure 3),

Figure 3: Established Japanese knotweed in bloom



several years of treatment may be required to fully exhaust the root reserves and eliminate new sprouts.

Management options should begin as soon as a new infestation is noted to improve efficacy and reduce the amount of time and materials required. Please contact your Bartlett Arborist Representative to learn about management strategies.



Founded in 1926, The Bartlett Tree Research Laboratories is the research wing of Bartlett Tree Experts. Scientists here develop guidelines for all of the Company's services. The Lab also houses a state-of-the-art plant diagnostic clinic and provides vital technical support to Bartlett arborists and field staff for the benefit of our clients.