

Keithia Blight

Bruce R. Fraedrich, PhD, Plant Pathologist

Keithia blight, a disease caused by the fungus *Didymascella*, leads to blighting, twig dieback, and branch death on species of *Thuja* and *Juniperus* (Figure 1). The pathogen is most damaging on western red cedar, *Thuja plicata*. The cultivars ‘Atrovirens’ and ‘Excelsa’ are the most commonly and severely affected.

Range

This disease is found throughout North America but is most common in the Pacific Northwest and western Canada. Small trees, lower branches of larger trees, and hedges are most likely to be affected because shaded and crowded conditions impede drying of the foliage favoring infection.

Symptoms

Symptoms begin as yellowing or browning of the individual infected leaf scales, usually in sharp contrast to the adjacent healthy tissue. One to several dark brown or black fruiting structures develop on the infected scales in late spring and by fall, the infected scales turn completely brown. Diseased foliage will either turn gray or drop off. Fruiting structures (apothecia) embedded in the tissue are clearly visible, but may eventually fall out leaving characteristic holes or pits (Figure 2). Affected scales are usually scattered amongst healthy tissue because the fungus is unable to progress from one scale to the next. Infection is primarily restricted to the current year’s foliage.

Disease Cycle

The principle means of infection is the sexual spores (ascospores), which are forcibly discharged from the fruiting structures and spread by wind and rain. Spore production begins in late spring and reaches a peak

Figure 1: Keithia blight on juniper



Figure 2: Infected leaf scales (tan) exhibiting pit-like fruiting structures



between August and November. Infection is favored by conditions that cause foliage to remain moist for extended periods of time.

Management

Cultural practices, such as proper spacing at planting, will promote foliage-drying and reduce the likelihood of infection. Shading can increase the severity of this and other foliage diseases. Pruning any over-story or adjacent plants to improve light and air penetration to lower foliage will help suppress disease. On trees that have lost foliage or branches due to this disease, good cultural practices including fertilization based on soil analysis, proper mulching, and irrigation during dry periods in the growing season will aid recovery. Chemical control of this disease requires the proper application of preventive fungicides. Please contact your Bartlett Arborist Representative to learn about management strategies.



Founded in 1926, The Bartlett Tree Research Laboratories is the research wing of Bartlett Tree Experts. Scientists here develop guidelines for all of the Company's services. The Lab also houses a state-of-the-art plant diagnostic clinic and provides vital technical support to Bartlett arborists and field staff for the benefit of our clients.