

# RESEARCH LABORATORY TECHNICAL REPORT



## Fall Webworm

By The Bartlett Lab Staff  
Directed by Kelby Fite, PhD

The fall webworm, *Hyphantria cunea* (Drury), is a native pest of shade and ornamental trees throughout much of North America. It attacks approximately 100 species of deciduous trees and shrubs in the United States. Preferred hosts will vary from region to region, but may include cherry, persimmon, black walnut, hickory, pecan, sweetgum and ash. The insect is easily recognized by the silken webs (nests) (Figure 1), which larvae construct on branch terminals during mid-to-late summer. These webs serve as protective structures where the larvae live and feed on the enclosed foliage. Two or more feet of the branch terminal are often enveloped by the webs, which become filled with cast skins, excrement and pieces of dried leaves as the larvae feed and mature. The webs turn black as autumn progresses and persist on trees long into winter, thereby rendering infested trees extremely unsightly. In addition, because the fall webworm is a sun-loving insect that prefers open grown roadside or ornamental plantings, webs in even lightly infested trees are extremely conspicuous.

Populations of the fall webworm are usually kept in check by naturally occurring parasites and predators. Subsequently, the main impact of this insect is the reduction in aesthetic value of shade and ornamental plants due to the unsightly webs. Periodically, however, outbreaks will occur resulting in substantial defoliation of host plants. Successive years of defoliation will weaken trees and shrubs and predispose them to attack by wood boring insects and disease-causing organisms.

### Description

Larvae of the fall webworm are approximately one to 1¼ inches in length when fully grown (Figure 3). They range in color from yellow to green to brown and often bear a black stripe down the back and a yellowish stripe

**Figure 1: Fall webworm silken webs (nests)**



**Figure 2: Female moth with eggs; Photo courtesy of Lacy L. Hyche, Auburn University, Bugwood.org**



reddish tubercles (raised spots) from which issue long, light-colored hairs. Adult moths are stout-bodied with a wingspan of 1 ¼ to 1 ½ inches. They may be pure white or white with brown markings. Eggs are present in masses on the underside of host leaves. They are light green in color, but usually covered with white scales from the female moth.

**Figure 3: Fall webworm larva**



### Life Cycle

Moths are present from May to July depending on the region of the country. Following mating, eggs are laid in masses of 400-500 and hatch within seven to ten days. Young larvae immediately begin to construct webs, enveloping the foliage at the branch tips. As more food is needed by the developing larvae, webs are expanded to include more foliage. The larvae are gregarious and live together as a colony within the web until the last instar stage, at which time they begin to disperse in search of pupation sites. The insect overwinters as pupae within the duff or upper few inches of soil.

One generation per year occurs in the northern latitudes, while two or more generations occur farther south. A partial second generation may occur in the latitudes corresponding to southern New York and New Jersey. In the South, larvae of the first generation are present during June and early July. Second generation larvae and larvae in areas having only one generation per year are noticeable from mid-July until early October.

### Control

On small trees and shrubs, suitable control can be obtained by pruning out and destroying webs while they are still small. On large plantings, chemical control is necessary. Please contact your Bartlett Arborist Representative for treatment options.

Plants severely defoliated by the fall webworm should be fertilized in late fall or early spring in order to maintain vigor and offset the debilitating effects of premature foliage loss.



**Founded in 1926, The Bartlett Tree Research Laboratories is the research wing of Bartlett Tree Experts. Scientists here develop guidelines for all of the Company's services. The Lab also houses a state-of-the-art plant diagnostic clinic and provides vital technical support to Bartlett arborists and field staff for the benefit of our clients.**